

DECOMPOSITION MATRICES FOR EXCEPTIONAL GROUPS AT $d = 4$

OLIVIER DUDAS AND GUNTER MALLE

ABSTRACT. We determine the decomposition matrices of unipotent ℓ -blocks of defect Φ_4^2 for exceptional groups of Lie type up to a few unknowns. For this we employ the new cohomological methods of the first author, together with properties of generalized Gelfand-Graev characters which were recently shown to hold whenever the underlying characteristic is good.

1. INTRODUCTION

The main aim of this paper is to determine the decomposition matrices for the unipotent blocks of finite exceptional groups of Lie type $E_6(q)$, ${}^2E_6(q)$, $E_7(q)$, $E_8(q)$ and $F_4(q)$ for odd primes ℓ dividing $\Phi_4(q) = q^2 + 1$. For these groups, we study only the blocks with defect at most 2 — which amounts to excluding only the principal block of $E_8(q)$ — and we obtain an approximation to the decomposition matrices in that a small number of entries remain undetermined for ${}^2E_6(q)$ and $F_4(q)$. On the way, we also find decomposition matrices for orthogonal groups of rank up to 7 which occur as Levi subgroups. As a byproduct, we obtain the repartition of the simple unipotent modules into Harish-Chandra series.

The decomposition matrices are determined inductively, by the combination of standard methods like Harish-Chandra induction and restriction, and the new ingredient introduced in [9] to tackle the discrete series. Our strategy can be summarized in the following three steps.

- Step 1.* We start by using Harish-Chandra induction from proper Levi subgroups, Harish-Chandra restriction from suitable overgroups and decomposition numbers for Hecke algebras to compute the columns of the decomposition matrix corresponding to the non-cuspidal simple modules. In several cases this determines entirely the unipotent part of the decomposition matrix.
- Step 2.* We consider suitable generalized Gelfand–Graev characters containing the missing columns. The properties of these projective characters (see [20, 23]) force the decomposition matrix to be unitriangular, but their construction introduce some conditions on the underlying characteristic of the groups considered.
- Step 3.* Finally, we use virtual projective characters afforded by cohomology of Deligne–Lusztig varieties. As observed in [9], we have some control on the multiplicity of the various PIMs in these virtual characters, from which we deduce upper bounds on the

Date: November 5, 2014.

2010 Mathematics Subject Classification. Primary 20C33; Secondary 20G40.

The second author gratefully acknowledges financial support by ERC Advanced Grant 291512.

missing decomposition numbers. In many cases these bounds are small enough to determine the numbers.

The paper is built up as follows. In Section 2 we present the general methods used in many of the arguments. Then in Section 3 we determine decomposition matrices for some orthogonal groups of type D_n , $n \leq 8$. In the following sections, we consider the exceptional groups of type E_n , $6 \leq n \leq 8$. For these we are able to determine all the decomposition numbers for blocks of defect at most 2, which excludes only the principal block of groups of types D_7 , D_8 and E_8 . In Section 7 we turn to the twisted groups of types 2D_n , $n \leq 7$, and 2E_6 . We finish by the case of symplectic groups of type C_n , $n \leq 4$ and exceptional groups of type F_4 in Section 8. In those cases we have to assume that the underlying characteristic is good, and even then we are not able to determine all decomposition numbers. Some entries remain unknown, but we still give conditions and relations that they satisfy.

Let us note that our calculations give rather large examples for Geck's conjectures on the shape of ℓ -decomposition matrices (see [12, Conjecture 3.4] for a precise formulation of the conjectures). Indeed, in all the cases we consider we observe that:

- the decomposition matrix has a unitriangular shape, with respect to an order compatible with Lusztig's a -function,
- within a given family, the square submatrix is the identity matrix (up to some indeterminates for types D_7 , 2D_7 , 2E_6 and C_4),
- any cuspidal unipotent character remains irreducible after ℓ -reduction (more generally any unipotent character with smallest a -function within its Harish-Chandra series).

2. METHODS

We determine decomposition matrices for unipotent blocks of various families of groups of Lie type G , where $G = \mathbf{G}(q)$ is the group of fixed points under a Frobenius endomorphism with respect to an \mathbb{F}_q -structure of a simple algebraic group \mathbf{G} over the algebraic closure of \mathbb{F}_q . More precisely, we consider the case that ℓ is an odd prime dividing $q^2 + 1$. In particular, we have $\ell \geq 5$ always. In the proofs we make use of several standard arguments which we collect here for easier reference.

Firstly, the subdivision of unipotent characters into ℓ -blocks is known in our situation, see [3]. Secondly, by results of Geck and Hiss, whenever ℓ is a good prime for the group in question, then the unipotent characters form a basic set for the union of unipotent blocks. Thus, for good primes the decomposition matrix for a unipotent block is known once the decomposition numbers for the unipotent characters in that block have been found. To determine the decomposition matrix of the block is hence equivalent to finding the (unipotent parts of the) ordinary characters of all projective indecomposable modules (PIMs) in that block.

One standard method for constructing projective characters is via Harish-Chandra induction R_L^G of projective characters from proper Levi subgroups L , which we may assume to be known by induction. Thus our first source for projective characters is

(HCi) Harish-Chandra induction of projective characters from proper Levi subgroups.

This Harish-Chandra induction can be computed explicitly in terms of induction in relative Weyl groups. All of our calculations were done in the **Chevie**-system [21]. In addition, Harish-Chandra restriction ${}^*R_L^G$ of projective characters also yields projective characters. This leads to the following indecomposability criterion:

(HCr) Let χ be a projective character of G . If no proper subcharacter of χ has the property that its Harish-Chandra restriction to Levi subgroups L decomposes non-negatively on the PIMs of L , then χ is the character of a PIM.

One of our results is the subdivision into modular Harish-Chandra series of the Brauer characters in the block. A valuable criterion to determine this is given by [13, Thm. 4.2]:

(Csp) The group G has a cuspidal unipotent Brauer character if and only if a Sylow ℓ -subgroup of G is not contained in any proper Levi subgroup of G .

Furthermore, the ordinary Gelfand–Graev character always provides the Steinberg PIM:

(St) There exists a PIM with unipotent part just the ordinary Steinberg character. It is non-cuspidal if and only if a Sylow ℓ -subgroup of G is contained in a proper Levi subgroup L . In this case it is a summand of the Harish-Chandra induction of the Steinberg PIM from L .

Other PIMs will appear as direct summands of generalized Gelfand–Graev representations (GGGRs). The results of Lusztig [20, §11], which have recently been extended to good characteristic by Taylor (see [23, Thm. 14.10]), give an approximation of some columns of the decomposition matrix:

(GGGR) Assume the underlying characteristic of G is good. Given a unipotent character ρ , there exists a (projective) Gelfand–Graev representation Γ such that ρ occurs in Γ , and any other unipotent constituent in Γ is either in the same family as ρ or has a larger a -value.

Under suitable conditions on the family of unipotent characters containing ρ one can even use [5, Thm. 6.5(ii)] to compute the multiplicities of the characters in the family in a GGGR. An instructive example is given in the proof of Theorem 3.6.

A further tool is given by a particular case of Dipper’s result (see [6, 4.10] for the precise assumptions):

(End) The decomposition matrix of the Hecke algebra $\text{End}_G(R_T^G(\mathbb{Z}_\ell))$ embeds as a submatrix into the decomposition matrix of G .

Dipper’s result holds more generally for a Hecke algebra associated with $R_L^G(\rho)$ where ρ is a cuspidal unipotent character satisfying the following two conditions (see [11, §2.6]):

- the ℓ -reduction of ρ is an irreducible Brauer character φ ,
- $N_W(W_L, \rho) = N_W(W_L, \varphi)$

where W denotes the Weyl group of G . Note that when L is classical, ρ is the unique cuspidal unipotent character, so that $N_W(W_L, \rho) = N_W(W_L)$ and the second condition is automatically satisfied. The first condition is conjectured to hold whenever ℓ is good, as already mentioned in the introduction.

For dealing with PIMs which are not induced from a proper Levi subgroup, we will make use of suitably chosen Deligne–Lusztig characters as in [9] and [7, Sec. 6]:

(DL) Let χ be a projective character of G . If $w \in W$ is minimal in the Bruhat order for the property that the unipotent part of χ occurs in the Deligne–Lusztig character R_w , then the sign of its multiplicity in R_w is $(-1)^{\ell(w)}$.

We shall often use the following particular case:

(Cox) Let χ be the character of the projective cover of a cuspidal unipotent module, and $w \in W$ be a Coxeter element. Then the multiplicity of the unipotent part of χ in $(-1)^{\ell(w)} R_w$ is non-negative.

The previous two arguments will usually give upper bounds on decomposition numbers. Lower bounds can be obtained from ℓ -reduction of non-unipotent characters, which are non-negative combinations of irreducible Brauer characters. This applies in particular to the Deligne–Lusztig induction of characters in general position:

(Red) Let T_w be a torus of type $w \in W$. Assume that there exists an ℓ -character of T_w in general position. Then the ℓ -reduction of $(-1)^{\ell(w)} R_w$ is a non-negative combination of irreducible Brauer characters.

Indeed, if θ is an ℓ -character of T_w in general position then $(-1)^{\ell(w)} R_{T_w}^G(\theta)$ is an irreducible character by [4] and it has the same ℓ -reduction as $(-1)^{\ell(w)} R_w = (-1)^{\ell(w)} R_{T_w}^G(1_{T_w})$.

We will also use variations of the following elementary observation:

(Sum) Let $\chi_1 + \chi_2, \chi_1 + \chi_3, \chi_2, \chi_3$ be characters of projectives modules, and assume that χ_2, χ_3 are indecomposable. Then χ_1 is the character of a projective module.

Indeed, we have two direct sum decompositions of a projective module with character $\chi_1 + \chi_2 + \chi_3$, and the theorem of Krull–Schmidt allows to conclude. Sometimes, we will also make use of the following obvious fact:

(Deg) The degrees of the irreducible Brauer characters in a block can be computed from the inverse of the decomposition matrix; they are all positive.

Our notation for modular Harish-Chandra series is as follows: characters in the principal series are labelled "ps", or sometimes "p" for short in large tables. If a Levi subgroup has a single cuspidal Brauer character, its Harish-Chandra series is labelled by the Dynkin type of that Levi subgroup. Else, it is labelled by the name of the corresponding ordinary unipotent character. For these, in turn, as customary we use the labelling in terms of ordinary Harish-Chandra series. Cuspidal Brauer characters are labelled by "c".

3. DECOMPOSITION MATRICES FOR ORTHOGONAL GROUPS OF TYPE D_n

We first determine the decomposition matrices of orthogonal groups $\mathrm{SO}_{2n}^+(q)$, with $n \leq 7$ and q a prime power, for primes $2 \neq \ell \mid (q^2 + 1)$, except in the case that $(q^2 + 1)_\ell = 5$. In the latter case, the decomposition matrices can be expected to be different from those in the general case, see Remarks 3.2 and 3.4.

3.1. Decomposition matrices for $\mathrm{SO}_8^+(q)$. Let first $n = 4$, so $G = \mathrm{SO}_8^+(q)$. For of the unipotent characters of G lie in ℓ -blocks of defect zero, the others lie in the principal block. Miyachi [22, Lemma 9] gives an approximation of the Φ_4 -modular decomposition matrix of the principal block of $\mathrm{SO}_8^+(q)$ when q is odd, based on the triangularity of the decomposition matrix proved by Geck–Pfeiffer [16] using generalized Gelfand–Graev characters. Here we extend their results to all q and determine the missing entry.

TABLE 1. $\mathrm{SO}_8^+(q)$, $(q^2 + 1)_\ell > 5$

Here, D_3, A_3, A'_3 denote three non-conjugate Levi subgroups of type A_3 .

Finally, we use (Cox) to determine a : the generalized 1-eigenspace of the Frobenius endomorphism on the Deligne–Lusztig character associated with a Coxeter element decomposes as

Remark 3.2. The 5-modular decomposition matrices of $\mathrm{SO}_8^+(2)$ and $\mathrm{SO}_8^+(3)$ are known; they differ from the one in Table 1 in that the entry “2” is replaced by “1”. Thus, Theorem 3.1 does not extend to the case $(q^2 + 1)_\ell = 5$.

Theorem 3.3. *Let ℓ be a prime. The ℓ -modular decomposition matrices for the unipotent blocks of $\mathrm{SO}_{10}^+(q)$ of positive defect for $(q^2 + 1)_\ell > 5$ are as given in Tables 2 and 3.*

TABLE 2. $\mathrm{SO}_{10}^+(q)$, $(q^2 + 1)_\ell > 5$ [illegible]

TABLE 3. $\mathrm{SO}_{10}^+(q)$, block of defect 1, $2 \neq \ell | (q^2 + 1)$

$$.41 \text{ --- } 2.21 \text{ --- } 1^2.21 \text{ --- } .21^3 \text{ --- } \bigcirc$$

$ps \qquad ps \qquad ps \qquad D_3$

Using (HCi) gives $\tilde{\Psi}_i$ for $i \in \{2, 3, 4, 5, 8, 10, 12\}$. The decomposition matrix of the Hecke algebra of type D_5 at a fourth root of unity gives by (End) the seven principal series PIMs Ψ_1, \dots, Ψ_5 , Ψ_7 and Ψ_{10} for all primes ℓ with $(q^2 + 1)_\ell > 5$. Furthermore, (HCi) yields $\tilde{\Psi}_9 := \Psi_9 + \Psi_8$ and $\tilde{\Psi}_{10} := \Psi_9 + \Psi_{10}$. An application of (Sum) yields Ψ_9 .

The centralizer of a Sylow Φ_4 -torus of G is contained in a Levi subgroup L of type D_4 , so [13, Thm. 4.2] shows that the Harish-Chandra induction $\tilde{\Psi}_{13} = \Psi_{13} + \Psi_{14}$ of the Steinberg PIM from L has two summands, namely Ψ_{13} and Ψ_{14} .

Finally, using (HCr) we check that, independently from the value of a , all Ψ_i are indecomposable. Indeed, no proper subsums restrict to a non-negative combination of PIMs in all Levi subgroups. We shall prove that $a = 2$ in Theorem 3.6 below. \square

3.3. Decomposition matrices for $\mathrm{SO}_{12}^+(q)$. Now let $G = \mathrm{SO}_{12}^+(q)$. This group has four unipotent Φ_4 -blocks of defect zero and three blocks of defect Φ_4^2 . We label these blocks by the symbol for their 4-Harish-Chandra source in a 4-split Levi subgroup of type $\mathrm{SO}_4^+(q)$.

TABLE 4. $\mathrm{SO}_{12}^+(q)$, block $\binom{2}{0}$, and $\mathrm{SO}_{16}^+(q)$, block $\binom{3}{1}$, $(q^2 + 1)_\ell > 5$

[illegible]

Proof. All projectives listed in the tables are obtained by (HCi) from the Levi subgroups of types D_5 and A_5 , except that instead of Ψ_4 and Ψ_{11} in the first block we obtain $\Psi_4 - (2 - a)(\Psi_{12} - \Psi_{14})$ and $\Psi_{11} + (2 - a)(\Psi_{12} - \Psi_{14})$, and instead of Ψ_5 and Ψ_{12} in the third block we obtain $\Psi_5 - (2 - a)(\Psi_{11} - \Psi_{14})$ and $\Psi_{12} + (2 - a)(\Psi_{11} - \Psi_{14})$, with the parameter $a \in \{0, 1, 2\}$ as in the proof of Theorem 3.3. It is straightforward to check by

[illegible][illegible]

3.4. Decomposition matrices for $\mathrm{SO}_{14}^+(q)$. We now consider the three blocks of positive Φ_4 -defect for the groups $G = \mathrm{SO}_{14}^+(q)$. The non-principal block is again labelled by the symbol for its 4-Harish-Chandra source in a 4-split Levi subgroup of type $\mathrm{SO}_6^+(q)$.

(a) *If q is odd then the decomposition matrix for the principal ℓ -block of G is as given in Tables 7 and 8.*

- (b) *The decomposition matrix for the non-principal unipotent ℓ -block G of positive defect is as given in Table 9.*

Proof. Let us first consider the block with defect Φ_4^2 . We argue how to construct projectives Ψ_1, \dots, Ψ_{14} with unipotent part equal to the columns in Table 9. (HCi) and (End) give all Ψ_i , except that instead of Ψ_4 and Ψ_{13} we find $\Psi_4 - (2-a)(\Psi_{10} - \Psi_{14})$ and $\Psi_{13} + (2-a)(\Psi_{10} - \Psi_{14})$, with $a \in \{1, 2\}$ as in the proof of Theorem 3.5. Again, it is easily seen by (HCr) that all these characters are indecomposable. Now the tenth Brauer character has positive degree only if $a \geq 2$, which shows by (Deg) that $a = 2$ in this table and also in the decomposition matrices for $\mathrm{SO}_{10}^+(q)$ and $\mathrm{SO}_{12}^+(q)$, thus completing the proofs of Theorems 3.3 and 3.5. Note that up to this point we did not use any assumption on q .

We now turn to the principal block. Here, (HCi) yields Ψ_i except for $i \in \{2, 14, 16, 19, 24, 25, 26, 34, 35, 36, 38, 40\}$. Using (Sum) we get other projective characters: $\Psi_2 + \Psi_3$ and $\Psi_2 + \Psi_4 + \Psi_{10}$ give Ψ_2 , $\Psi_{14} + \Psi_{17}$ and $\Psi_{14} + \Psi_{29}$ give Ψ_{14} , $\Psi_6 + \Psi_{19}$ and $\Psi_{19} + \Psi_{30}$ give Ψ_{19} , $\Psi_{12} + \Psi_{16}$ and $\Psi_{16} + \Psi_{19}$ give Ψ_{16} , $\Psi_9 + \Psi_{20} + \Psi_{24}$ and $\Psi_{24} + \Psi_{27}$ give Ψ_{24} , $\Psi_{31} + \Psi_{35}$ and $\Psi_{33} + \Psi_{35}$ give Ψ_{35} , $\Psi_{28} + \Psi_{34}$ and $\Psi_{34} + \Psi_{35}$ give Ψ_{34} . Furthermore we find Ψ_{26} with $c = 1$. The PIM Ψ_{40} is cuspidal by (St). (HCr) shows that all of the projectives obtained so far, with the possible exception of Ψ_{26} , which might contain Ψ_{27} once, are indecomposable. We have thus obtained all but three columns of the decomposition matrix. Since we have accounted for all proper Harish-Chandra series, the remaining three Brauer characters must be cuspidal.

To establish the unitriangularity we look at suitably chosen generalized Gelfand–Graev representations and use (GGGR). Note that we have to assume that q is odd in order to construct these representations and use the results in [23]. Let us first consider the family $\mathcal{F} = \{\rho_{1^5.2}, \rho_{1.21^4}, \rho_{2.21^3}, \rho_{D_4.1^3}\}$ of unipotent characters. The special character of this family is $\rho_{1.21^4}$; via the Springer correspondence, it corresponds to a special unipotent class, and we denote by \mathcal{O} its dual. By [23, Thm. 14.10], the character of any GGGR attached to \mathcal{O} involves characters lying in \mathcal{F} or in a family with a strictly larger a -value than that of \mathcal{F} . In particular, the only characters in the block that can occur are $\rho_{1^5.2}, \rho_{1.21^4}, \rho_{D_4.1^3}, \rho_{1^3.1^4}, \rho_{1^2.1^5}, \rho_{1.1^6}$ and $\rho_{1.1^7}$, which gives an approximation of Ψ_{36} . Furthermore, $u \in \mathcal{O}^F$ satisfies the following two conditions:

- the small finite group attached to the family as in [19, Chap. 4] and the component group $A_G(u) := (C_G(u)/C_G(u)^\circ)^F$ are isomorphic (to $\mathbb{Z}/2\mathbb{Z}$),
- at most one of the local systems on (u) is not in the principal block,

in which case one can apply [5, Thm. 6.5(ii)] to compute the projection of the GGGR Γ_u into the span of \mathcal{F} . Recall that the characters in a family are parametrized by pairs (g, ψ) where g runs over a set of representatives of conjugacy classes of the small finite group attached to the family, say \mathcal{G} and $\psi \in \mathrm{Irr}(C_{\mathcal{G}}(g))$. For $g \in \mathcal{G}$, the Mellin transform of the pair $(g, 1)$ is given by

$$\mu_{(g,1)} = \sum_{\psi \in \mathrm{Irr}(C_{\mathcal{G}}(g))} \psi(g) \rho_{(g,\psi)}.$$

TABLE 7. $\mathrm{SO}_{14}^+(q)$, q odd, principal block, $(q^2 + 1)_\ell > 5$

[illegible]

[illegible]

The unipotent part of the Gelfand-Graev representation of $\mathrm{SO}_{16}^+(q)$ associated with the family $\{\rho_{12,16}\}$, cut by the principal block, is of the form $\rho_{12,16} + \alpha\rho_{2,16} + \beta\rho_{1,8}$ for

projective characters, each of which contains only one unipotent character of the family. If Kawanaka's conjecture holds for the family $\{1^2.2^21, 1^3.2^2, 1.2^3, D_4 : .3\}$ of $\mathrm{SO}_{14}(q)$, then b_1 and d must be equal to zero. More generally, as suggested by Geck, Kawanaka's characters should force block unitriangularity of the decomposition matrix whenever p is good.

Our method is not sufficient to determine all the decomposition numbers of $\mathrm{SO}_{14}^+(q)$. However, we can use [8, Conj. 1.2] to determine small upper bounds for the missing entries. Following [8], we denote by Q_w the virtual character afforded by the Alvis–Curtis dual of the intersection cohomology of the Deligne–Lusztig variety corresponding to w . In addition, if $\lambda \in \overline{\mathbb{F}}_\ell$ we consider the virtual character $Q_w[\lambda]$ afforded by the generalized λ -eigenspace of the Frobenius on the intersection cohomology. Up to a sign, $Q_w[\lambda]$ is a proper character and Conjecture 1.2 in [8] predicts that it is actually the unipotent part of a projective character. The multiplicities of the various PIMs in $Q_w[\lambda]$ depend on the decomposition numbers (including the missing entries), forcing some linear combinations of decomposition numbers to be non-negative.

Proposition 3.8. *Assume Conjecture 1.2 in [8] holds. Then in the decomposition matrix of the principal Φ_4 -block of $\mathrm{SO}_{14}^+(q)$, we have $b_1 = b_3 = b_4 = b_5 = b_8 = 0$, $b_6, b_7, b_{10}, b_{11} \leq 2$, $b_9 \leq 6$, $b_{12}, b_{13} \leq 12$, $b_{14} \leq 18$ and $b_{15} \leq 20$.*

Proof. To obtain the upper bounds on the b_i 's we consider the characters $Q_{w_1}[q^3]$ for $w_1 = s_1 s_2 s_3 s_1 s_2 s_3 s_5 s_4 s_3 s_6 s_5 s_4 s_3 s_7 s_6 s_5 s_4 s_3$ and $Q_{w_2}[1]$ for $w_2 = s_1 s_2 s_3 s_1 s_2 s_3 s_5 s_6 s_5 s_7 s_6 s_5 s_4$ as well as their decomposition on the basis of PIMs.

The coefficient of Ψ_{26} in $Q_{w_1}[q^3]$ is $10 - 14b_1$. By [8, Conj. 1.2] it must be non-negative, which forces $b_1 = 0$. The list of coefficients of $\Psi_{27}, \dots, \Psi_{32}$ on $Q_{w_2}[1]$ is given by $-7b_3$, $-7b_4$, $-7b_5$, $17 + 7b_4 - 7b_6$, $17 - 7b_7$ and $7b_4 - 7b_8$. Since they must be all non-negative, we get $b_3 = b_4 = b_5 = b_8 = 0$ and $b_6, b_7 \leq 2$. With these values, the coefficients of $\Psi_{33}, \dots, \Psi_{40}$ on $Q_{w_2}[1]$ are

$$\begin{aligned} &14 + 7b_6 + 7b_7 - 7b_9, \quad 7b_7 - 7b_{10}, \quad 19 - 7b_{11}, \\ &62 + 7b_6 + 7b_7 - 7b_{12}, \quad -9 - 7b_6 - 7b_7 + 7b_9 + 7b_{12} - 7b_{13}, \\ &60 - 7b_6 - 7b_7 + 7b_{10} + 7b_{12} - 7b_{14} + d(9 + 7b_6 + 7b_7 - 7b_9 - 7b_{12} + 7b_{13}), \\ &42 + 21b_6 + 21b_7 - 21b_9 - 7b_{10} + 14b_{11} - 21b_{12} + 21b_{13} + 7b_{14} - 7b_{15}. \end{aligned}$$

From the first line we deduce $b_9 \leq b_6 + b_7 + 2$, $b_{10} \leq b_7$ and $b_{11} \leq 2$. From the second, $b_{12} \leq b_6 + b_7 + 8$ and $b_{13} \leq -2 - b_6 - b_7 + b_9 + b_{12}$, so that $b_{13} \leq b_{12}$. Finally, the last two lines yield $b_{14} \leq 8 - b_6 - b_7 + b_{10} + b_{12} \leq 16 + b_{10}$ and $b_{15} \leq 6 + 3b_6 + 3b_7 - 3b_9 - b_{10} + 2b_{11} - 3b_{12} + 3b_{13} + b_{14} \leq 16 + 2b_{11}$. \square

3.5. Decomposition matrices for $\mathrm{SO}_{16}^+(q)$. Finally, we consider the three blocks of Φ_4 -defect Φ_4^2 for the groups $G = \mathrm{SO}_{16}^+(q)$. They are again labelled by the symbol for their 4-Harish-Chandra source in a 4-split Levi subgroup of type $\mathrm{SO}_8^+(q)$.

Theorem 3.9. *The ℓ -modular decomposition matrices for the unipotent blocks of $\mathrm{SO}_{16}^+(q)$ of defect Φ_4^2 , for primes ℓ with $(q^2 + 1)_\ell > 5$, are as given in Tables 4, 6, 10 and 11.*

The two blocks of $\mathrm{SO}_{12}^+(q)$ and $\mathrm{SO}_{16}^+(q)$ in Table 4 are Morita equivalent, as well as the two blocks in Table 6.

TABLE 10. $\mathrm{SO}_{16}^+(q)$, block $\begin{pmatrix} 12 \\ 03 \end{pmatrix}$, $(q^2 + 1)_\ell > 5$

[illegible]

Here, all character degrees have been divided by $q^3\Phi_4^2\Phi_7\Phi_{12}$.

TABLE 11. $\mathrm{SO}_{16}^+(q)$, block $\begin{pmatrix} 2 & 3 \\ 0 & 1 \end{pmatrix}$, $(q^2 + 1)_\ell > 5$

[illegible]

Here, all character degrees have been divided by $q^3\Phi_4^2\Phi_{12}\Phi_{14}$.

Proof. Harish-Chandra induction sends the fourteen PIMs in the principal block of $\mathrm{SO}_{12}^+(q)$ to the fourteen listed projectives of the first block of $\mathrm{SO}_{16}^+(q)$ in Table 4, and it sends irreducible characters to irreducible characters. Thus, by [2, Thm. 0.2] those two blocks are

Morita equivalent. Exactly the same assertions hold for the third block of $\mathrm{SO}_{12}^+(q)$ and the fourth block of $\mathrm{SO}_{16}^+(q)$ in Table 6.

In the second block of G , (HCi) yields all columns in Table 10, except for the second one. This is then obtained from the projectives $\Psi_1 + \Psi_2$ and $\Psi_2 + \Psi_4$ via (Sum).

In the third block, we obtain all Ψ_i in Table 11 for $i \notin \{1, 5, 6, 9\}$. Then, using (Sum) $\Psi_4 + \Psi_6$ and $\Psi_6 + \Psi_7$ give Ψ_6 , $\Psi_2 + \Psi_9$ and $\Psi_7 + \Psi_9 + \Psi_{11}$ give Ψ_9 , $\Psi_4 + \Psi_5$ and $\Psi_5 + \Psi_9$ give Ψ_5 , and finally $\Psi_1 + \Psi_2$ and $\Psi_1 + \Psi_5$ give Ψ_1 . \square

4. DECOMPOSITION MATRICES FOR $E_6(q)$

We now turn to decomposition matrices for the exceptional Lie type groups. We first consider $G = E_6(q)$ for primes $\ell \nmid \Phi_4(q)$ in which case the Sylow ℓ -subgroups are abelian homocyclic of rank 2. Note that again we do not need and will not specify the isogeny type of G , since the decomposition numbers of the unipotent characters will not depend on such a choice. The group $E_6(q)$ has ten unipotent blocks of ℓ -defect zero, one of cyclic defect and the principal block containing 16 unipotent characters.

Here, the decomposition matrix of the principal block has been determined by Miyachi [22, Thm. 37] except for three missing entries, which coincide with entries of the decomposition matrix for $D_4(q)$, again under the assumption that q is a power of a good prime. We give an independent proof of his result, valid for all prime powers q , and find the remaining entries using Theorem 3.1:

Proposition 4.1. *Let ℓ be a prime. Then the ℓ -modular decomposition matrices for the unipotent blocks of $E_6(q)$ of positive defect, for $(q^2 + 1)_\ell > 5$, are as given in Tables 12 and 13. In particular, the three undetermined entries in the ℓ -modular decomposition matrix of $E_6(q)$ in [22, Thm. 37] are all equal to 2.*

Proof. The Brauer tree for the block with cyclic defect is easily determined. We will construct PIMs Ψ_1, \dots, Ψ_{16} with unipotent parts as given by the columns of Table 12.

The PIMs in the principal series can be read off from the Φ_4 -modular decomposition matrix of the Iwahori–Hecke algebra of type E_6 given in [14, Tab. 7.13]. Note that by [15, Thm. 3.10] this agrees with the ℓ -modular decomposition matrix whenever $\ell \geq 5$ (since 5 is a good prime for E_6 and 20 does not divide any degree of E_6). So we have columns i in Table 12 for $i \in \{1, 2, 3, 4, 6, 7, 8, 9\}$. Furthermore, Ψ_{11} and Ψ_{13} are Harish-Chandra induced from Levi subgroups of types D_5 and A_5 .

The Hecke algebra for the ordinary cuspidal unipotent character of D_4 is of type A_2 with parameter q , thus remains semisimple modulo ℓ . Now Harish-Chandra induction from the Levi subgroup L of type D_5 yields projective characters $\Psi_5 + \Psi_{10}$ and $\Psi_{10} + \Psi_{14}$. Both of these must be the sum of two projective characters. Using (HCr) we find that $\Psi_5, \Psi_{10}, \Psi_{14}$ are the only subsums which can be projective.

Since all other Harish-Chandra series have been accounted for, and G cannot have cuspidal Brauer characters by (Csp), the missing three PIMs must lie in the series of the cuspidal Brauer character ρ_{14} of D_4 . Its relative Weyl group is the symmetric group \mathfrak{S}_3 , so the corresponding Hecke algebra must be semisimple. Now Harish-Chandra induction yields $\Psi_{12} + \Psi_{15}$ and $\Psi_{15} + \Psi_{16}$, and via (HCr) there is a unique way for each of these to split into sums of two non-zero projective characters. This completes the construction of Ψ_1, \dots, Ψ_{16} and thus the proof. \square

TABLE 12. $E_6(q)$, $(q^2 + 1)_\ell > 5$

$\phi_{1,0}$	1	1																	
$\phi_{6,1}$	$q\Phi_8\Phi_9$. 1																	
$\phi_{15,5}$	$\frac{1}{2}q^3\Phi_5\Phi_6^2\Phi_8\Phi_9$. . 1																	
$\phi_{15,4}$	$\frac{1}{2}q^3\Phi_5\Phi_8\Phi_9\Phi_{12}$	1 1 . 1																	
$D_4 : 3$	$\frac{1}{2}q^3\Phi_1^4\Phi_3^2\Phi_5\Phi_9$ 1																	
$\phi_{81,6}$	$q^6\Phi_3^3\Phi_6^2\Phi_9\Phi_{12}$. 1 1 . . 1																	
$\phi_{80,7}$	$\frac{1}{6}q^7\Phi_2^4\Phi_5\Phi_8\Phi_9\Phi_{12}$. 1 . 1 . 1 1																	
$\phi_{10,9}$	$\frac{1}{3}q^7\Phi_5\Phi_6^2\Phi_8\Phi_9\Phi_{12}$	1 . . 1 . . . 1																	
$\phi_{90,8}$	$\frac{1}{3}q^7\Phi_3^3\Phi_5\Phi_6^2\Phi_8\Phi_{12}$. . 1 . . 1 . . 1																	
$D_4 : 21$	$\frac{1}{2}q^7\Phi_1^4\Phi_3^2\Phi_5\Phi_8\Phi_9$ 1																	
$\phi_{81,10}$	$q^{10}\Phi_3^3\Phi_6^2\Phi_9\Phi_{12}$ 1 1 . 1 . 1																	
$\phi_{15,17}$	$\frac{1}{2}q^{15}\Phi_5\Phi_6^2\Phi_8\Phi_9$ 2 . . . 1 . 1 1																	
$\phi_{15,16}$	$\frac{1}{2}q^{15}\Phi_5\Phi_8\Phi_9\Phi_{12}$ 1 . . 1 1 1																	
$D_4 : 1^3$	$\frac{1}{2}q^{15}\Phi_1^4\Phi_3^2\Phi_5\Phi_9$ 1																	
$\phi_{6,25}$	$q^{25}\Phi_8\Phi_9$ 1 . . 2 1 . 1 . 1																	
$\phi_{1,36}$	q^{36} 1 1 2 . 1																	
		$ps \ ps \ ps \ ps \ D_4 \ ps \ ps \ ps \ ps \ D_4 \ A_3 \cdot 1^4 \ A_3 \ D_4 \cdot 1^4 \cdot 1^4$																	

TABLE 13. $E_6(q)$, block of defect 1, $2 \neq \ell | (q^2 + 1)$

$$\phi_{20,2} \xrightarrow{ps} \phi_{60,5} \xrightarrow{ps} \phi_{60,11} \xrightarrow{ps} \phi_{20,20} \xrightarrow{A_3} \bigcirc$$

5. DECOMPOSITION MATRICES FOR $E_7(q)$

We next consider the four unipotent Φ_4 -blocks of $E_7(q)$ of positive defect (see [3, Tab. 2]), which we name by their 4-Harish-Chandra sources in a Levi subgroup of type $A_1(q)^3$.

Theorem 5.1. *The ℓ -modular decomposition matrices for the unipotent blocks of $E_7(q)$ of positive defect for primes ℓ with $(q^2 + 1)_\ell > 5$ are as given in Tables 14–17.*

Proof. For the principal block, all columns but the sixth are obtained by (HCi). The projectives $\Psi_6 + \Psi_7$ and $\Psi_6 + \Psi_{10}$ then yield Ψ_6 via (Sum).

For the second block, all but the ninth column are gotten by (HCi) and then $\Psi_9 + \Psi_{10}$ and $\Psi_9 + \Psi_{12}$ give Ψ_9 .

For the third block, all PIMs are Harish-Chandra induced. Finally, all projectives in the fourth block except for the fifth come from (HCi). The projectives $\Psi_5 + \Psi_7$ and $\Psi_5 + \Psi_{10}$ then give Ψ_5 via (Sum). Then (HCr) shows that all of these projectives are indecomposable. \square

TABLE 16. $E_7(q)$, block $2 \otimes 1^2 \otimes 1^2$, $(q^2 + 1)_\ell > 5$

[illegible]

TABLE 17. $E_7(q)$, block $1^2 \otimes 1^2 \otimes 1^2$, $(q^2 + 1)_\ell > 5$

[illegible]

ℓ -blocks of Φ_4 -defect two, see [3, Tab. 2]. These are labelled by the four Φ_4 -cuspidal unipotent characters of the Φ_4 -split Levi subgroup of type D_4 .

Theorem 6.1. *The ℓ -modular decomposition matrices for the unipotent blocks of $E_8(q)$ of defect Φ_4^2 for primes $\ell > 5$ dividing $q^2 + 1$ are as given in Tables 18–21.*

TABLE 20. $E_8(q)$, block $\binom{23}{01}$, $5 < \ell | (q^2 + 1)$

$\phi_{28,8}$	1																		
$\phi_{160,7}$.	1																	
$\phi_{300,8}$.	.	1																
$\phi_{972,12}$.	1	1	1															
$\phi_{840,14}$.	.	1	1	1														
$\phi_{700,16}$	1	1	.	.	.	1													
$D_4 : \phi_{12,4}$	1												
$D_4 : \phi'_{6,6}$	1											
$D_4 : \phi''_{6,6}$	1										
$\phi_{1344,19}$.	1	.	1	.	1	.	.	.	1									
$\phi_{840,26}$	1	1	1								
$\phi_{700,28}$.	.	.	1	1	1	.	1							
$\phi_{972,32}$	1	.	.	.	1	1	.	1						
$\phi_{300,44}$	2	.	1	.	1	1					
$\phi_{160,55}$	2	.	.	1	.	1	1	.	1				
$\phi_{28,68}$	1	.	.	2	.	.	.	1	.	.	.	1			
		ps	ps	ps	ps	ps	ps	D_4	D_4	D_4	ps	ps	A_3	A_3	$.1^4$	$.1^4$	$.1^4$		

TABLE 21. $E_8(q)$, block $\begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 3 \end{pmatrix}$, $5 < \ell | (q^2 + 1)$

$\phi_{56,19}$		1																	
$\phi_{1400,7}$.	1																
$\phi_{1008,9}$.	.	1															
$\phi_{3240,9}$.	1	1	1														
$\phi_{2240,10}$.	1	.	1	1													
$D_4 : \phi'_{4,7}$		1												
$\phi_{4200,15}$.	.	1	1	.	.	1											
$\phi_{3200,22}$		1	1	1										
$D_4 : \phi'_{8,9}$		1									
$\phi_{4536,23}$.	.	.	1	.	.	1	1	.	1								
$\phi_{1400,29}$.	.	.	1	1	1	1							
$\phi_{840,31}$		1	2	.	1	.	.	.	1						
$\phi_{1344,38}$		2	.	1	.	1	.	1	1					
$D_4 : \phi_{4,13}$		1				
$\phi_{560,47}$		2	1	1	.	1	.	1			
$\phi_{8,91}$		1	1	.	.	2	.	1		
		<hr/>																	
		ps	ps	ps	ps	ps	D_4	ps	ps	D_4	ps	A_3	$.1^4$	A_3	D_4	$.1^4$	$.1^4$		

$\Psi_9 + \Psi_{12}$ yield Ψ_9 , $\Psi_6 + 2\Psi_7 + \Psi_{10}$ and $\Psi_{10} + \Psi_{11}$ yield Ψ_{10} , and $\Psi_{11} + \Psi_{15}$ and $\Psi_{14} + \Psi_{15}$ yield Ψ_{15} .

In the block above $\begin{pmatrix} 12 \\ 03 \end{pmatrix}$, (HCi) yields all columns Ψ_i except for indices $i \in \{3, 6, 11\}$. Here, the projectives $\Psi_{11} + \Psi_{12}$ and $\Psi_{11} + \Psi_{14}$ yield Ψ_{11} , $\Psi_3 + \Psi_5 + \Psi_{11}$ and $2\Psi_3$ yield Ψ_3 , and $\Psi_6 + 2\Psi_9$ and $2\Psi_6$ yield Ψ_6 .

In the block above $\begin{pmatrix} 23 \\ 01 \end{pmatrix}$, (HCi) yields the columns Ψ_i with $i \in \{3, 4, 6, 11, 12, 13\}$. The information obtained in this way does not seem to yield strong enough conditions to determine the decomposition matrix completely. So here we use in addition the decomposition matrix of the Hecke algebra of type E_8 . In characteristic zero this can be found in [14, Tab. 7.15]. By [15, Thm. 3.10] this agrees with the decomposition matrix in characteristic ℓ for all $\ell \geq 7$. This yields in addition the principal series PIMs Ψ_i with $i \in \{1, 2, 5, 10\}$. To construct the missing projectives, let us first consider the characters in the Harish-Chandra series of the ordinary cuspidal character of a Levi subgroup of type D_4 . Here, the relative Weyl group W has type F_4 , and the Hecke algebra H has parameters q^4, q . But then all characters of H relevant for our block lie in semisimple blocks of H for all $\ell \geq 5$ by [1, Thm. 3.13]. Thus, the decomposition of induced projectives from a Levi subgroup of type E_7 can be read off from the character table of W . In the third block, we find $\Psi_7 + \Psi_8$, $\Psi_7 + \Psi_9$, which should have a projective summand in common. The only splitting of these projectives compatible with (HCr) is as given. This accounts for the projectives Ψ_7, Ψ_8, Ψ_9 in the D_4 -series. Finally, we consider the characters above the cuspidal character $.1^4$ of the Levi subgroup of type D_4 . Here again, the relative Weyl group has type F_4 ; the parameters of the corresponding Hecke algebra are already determined locally inside D_5 and D_4A_1 to be the same as for the ordinary cuspidal character considered above. So again by [1, Thm. 3.13], all characters in the $.1^4$ -Harish-Chandra series correspond to blocks of H of defect 0. Harish-Chandra induction from E_7 yields $\Psi_{14} + \Psi_{15}$ and $\Psi_{15} + \Psi_{16}$, with a common summand. Visible, the only possible splitting is as claimed.

Finally, in the block indexed by $\begin{pmatrix} 123 \\ 013 \end{pmatrix}$, (HCi) yields all columns Ψ_i except for $i \in \{2, 3, 7\}$. The projectives $\Psi_5 + \Psi_7 + \Psi_{10}$ and $2\Psi_7 + \Psi_8$ yield Ψ_7 , $\Psi_2 + \Psi_7$ and $2\Psi_2 + \Psi_4$ yield Ψ_2 , and $\Psi_1 + \Psi_3 + 2\Psi_7$ and $\Psi_2 + \Psi_3$ give Ψ_3 .

It is now a routine computation using (HCr) to check that none of the projectives constructed above can be decomposable. \square

6.1. The Φ_4 -blocks in untwisted groups. In the following table, we have collected some numerical information on the various Φ_4 -blocks of defect Φ_4^2 and Φ_4^3 whose decomposition matrices we have determined: the relative Weyl group $W_G(b)$ of the block (see [3, Tab. 1]) and the distribution of its Brauer characters into Harish-Chandra series:

Remark 6.2. (a) It emerges that the distribution into modular Harish-Chandra series in all examples considered only depends on the relative Weyl group.

(b) In addition to the first block of $\mathrm{SO}_{12}^+(q)$ and the first block of $\mathrm{SO}_{16}^+(q)$, and the third block of $\mathrm{SO}_{12}^+(q)$ and the fourth block of $\mathrm{SO}_{16}^+(q)$, which form Morita equivalent pairs by Theorem 3.9, the following four pairs of blocks have identical decomposition matrices (after suitably reordering the characters): the principal block of E_6 and the 3rd block of E_8 ; the 2nd and the 3rd block of E_7 ; the first blocks of E_7 and of E_8 ; the 4th blocks of E_7 and of E_8 . It is claimed (without proof) in [22, Rem. 34] that the first listed pair of blocks are in fact Morita equivalent. It would be interesting to see whether this is true for all pairs mentioned above.

TABLE 22. HC-series in Φ_4 -blocks of defect Φ_4^2 and Φ_4^3

G	b	$W_G(b)$	$ \text{IBr}(b) $	ps	A_3	D_3	D_4	$.1^4$	A_3D_3	c
D_4	2	$G(4, 2, 2)$	10	5	2	1	1	1		
D_6				5	2	1	1	1		
D_5	1, 3	$G(4, 1, 2)$	14	7	1	2	2	2		
D_6				7	1	2	2	2		
D_7				7	1	2	2	2		
D_8				7	1	2	2	2		
E_7				7		3	2	2		
E_6	1, 4	G_8	16	8		2	3	3		
E_7				8		2	3	3		
E_8				8		2	3	3		
D_7	1	$G(4, 1, 3)$	40	15	3	5	6	6	1	4

Note that the series A_3 and D_3 fuse in E_6 (and hence in E_7 and E_8).

(c) The decomposition matrix for $\text{SO}_8^+(q)$ and the one for the second block of $\text{SO}_{12}^+(q)$ have automorphisms induced by the non-trivial graph automorphisms of the underlying groups. But note that also the decomposition matrix for the principal block of E_6 has an automorphism fixing $\phi_{6,1}, \phi_{80,7}, D_4:2.1$ and $\phi_{6,25}$ and interchanging the other characters in pairs, and similarly, the decomposition matrix for the second block of E_7 has an automorphism of order two with fixed points $\phi_{280,8}$ and $\phi_{216,16}$.

7. DECOMPOSITION MATRICES FOR TWISTED TYPE GROUPS

We now turned to simply-laced groups of twisted type, viz. 2D_5 , 2D_6 and 2E_6 .

7.1. Decomposition matrices for $\text{SO}_{10}^-(q)$. The group $G = \text{SO}_{10}^-(q)$ has four unipotent ℓ -blocks for primes $2 \neq \ell | (q^2 + 1)$, the principal block, one block with cyclic defect and two blocks of defect zero.

Theorem 7.1. *Assume that q is odd. Then the ℓ -modular decomposition matrices for the unipotent blocks of $\text{SO}_{10}^-(q)$ of positive defect for primes ℓ with $(q^2 + 1)_\ell > 5$ are as given in Tables 23 and 24.*

Proof. The Brauer tree for the block with cyclic defect is easily seen to be as given in Table 24. We describe how to obtain projectives $\Psi_1, \dots, \Psi_{11}, \Psi_{13}, \Psi_{14}$ for the principal block as in Table 23.

Application of (HCi) yields Ψ_i with $i \in \{1, 3, 4, 5, 6, 7, 8, 9, 13\}$. Moreover, we obtain $\Psi_2 + \Psi_3, \Psi_2 + \Psi_6$, so that (Sum) gives Ψ_2 . (Alternatively, the projectives in the principal series are obtained from the Iwahori–Hecke algebra H of type B_4 , with parameters q^2 and q .) Next, (HCi) gives $\tilde{\Psi}_{10} = \Psi_{10} + \Psi_8$, $\tilde{\Psi}'_{10} = \Psi_{10} + \Psi_{11}$ and $\tilde{\Psi}_{11} = \Psi_{11} + \Psi_{13}$. Thus, $\tilde{\Psi}_{10} + \tilde{\Psi}_{11} = \tilde{\Psi}'_{10} + \Psi_8 + \Psi_{13}$, which shows that Ψ_8, Ψ_{13} occur as summands of $\tilde{\Psi}_{10} + \tilde{\Psi}_{11}$. Nonnegativity of decomposition numbers implies that Ψ_8 occurs in $\tilde{\Psi}_{10}$ and Ψ_{11} in $\tilde{\Psi}_{11}$,

TABLE 23. $\mathrm{SO}_{10}^{-}(q)$, q odd, $(q^2 + 1)_{\ell} > 5$

Here, $a \in \{0, 1\}$.

TABLE 24. $\mathrm{SO}_{10}^{-}(q)$, block of defect 1, $2 \neq \ell | (q^2 + 1)$

$$3.1 \text{ --- } 1.3 \text{ --- } \bigcirc \text{ --- } 1.1^3 \text{ --- } 1^3.1$$

$ps \quad \quad \quad {}^2D_2 \quad \quad {}^2D_2 \quad \quad \quad ps$

so we obtain Ψ_{10} and Ψ_{11} . Application of (HCr) also shows that Ψ_1, \dots, Ψ_{11} and Ψ_{13} are indecomposable. The last projective Ψ_{14} is given by (St).

The Hecke algebra for the cuspidal Brauer character of ${}^2D_2(q) \cong A_1(q^2)$ is of type B_3 , and the parameters are seen locally in the Levi subgroups of types 2D_4 and ${}^2D_3 \times A_1$ to be q^4 and q . From its decomposition matrix it follows that exactly eight simple modules lie in the corresponding Harish-Chandra series in G . Since the unipotent block with cyclic defect contains two of them, the principal block will contain the remaining six. We have then accounted for all non-cuspidal Harish-Chandra series, so the remaining Brauer character must all be cuspidal.

By (GGGR), the unipotent part of Ψ_{12} is $\rho_{\cdot, 2^2} + a\rho_{\cdot, 21^2} + b\rho_{\cdot, 1^4}$, where a and b are unknown. To compute b we use (Cox), which gives the relation $b \leq a + 2$. The relation $b \geq a + 2$ is obtained from the ℓ -reduction of the non-unipotent character obtained by Deligne–Lusztig induction of an ℓ -character in general position of a torus of order $(q + 1)(q^2 + 1)^2$ (such a character exists whenever $(q^2 + 1)_\ell > 5$). Finally, to obtain an upper bound for a , we consider the generalized Gelfand–Graev representations of $\mathrm{SO}_{14}^-(q)$ associated to the family $\{\rho_{1^5, 1}, \rho_{1^3, 1^3}, \rho_{1, 2^2, 1}, \rho_{\cdot, 32, 1}\}$, and more precisely the one whose projection to this family is $\rho_{\cdot, 32, 1} + \rho_{1, 2^2, 1}$. The character of this representation, cut by the block containing $\rho_{\cdot, 32, 1}$ is of the form $\rho_{\cdot, 32, 1} + \rho_{1, 2^2, 1} + \alpha\rho_{1, 1^5}$ by (GGGR). The Harish-Chandra restriction of this character yields $a \leq 1$. \square

Remark 7.3. The 5-modular decomposition matrix of $\mathrm{SO}_{10}^-(2)$ is known; there, the last three entries in the partially unknown 12th column of Table 23 read $(1, 0, 1)$, whence the case when $(q^2 + 1)_\ell = 5$ does behave differently.

Theorem 7.4. *Assume that q is odd. Then the ℓ -modular decomposition matrix for the non-principal unipotent block of $\mathrm{SO}_{14}^-(q)$ of defect Φ_4^2 , for $(q^2 + 1)_\ell > 5$, is as given in Table 25.*

	$\Phi_3\Phi_6$	1									
1.5	$\frac{1}{2}q^2\Phi_3\Phi_8\Phi_{10}\Phi_{14}$	1	1								
32.1	$\frac{1}{2}q^2\Phi_3\Phi_5\Phi_8\Phi_{14}$	1	.	1							
321.	$\frac{1}{2}q^2\Phi_2^4\Phi_6\Phi_{10}\Phi_{14}$.	.	.	1						
$31^2.1$	$q^4\Phi_3^2\Phi_6\Phi_8\Phi_{14}$.	.	.	1	1					
$2^2.1.1$	$q^5\Phi_3\Phi_5\Phi_6\Phi_{10}\Phi_{14}$.	.	1	.	.	1				
$1^3.3$	$q^7\Phi_5\Phi_8\Phi_{10}\Phi_{14}$	1	.	1			
1.32	$q^9\Phi_3\Phi_5\Phi_6\Phi_{10}\Phi_{14}$	1	1	1	1		
3.1^3	$q^{10}\Phi_5\Phi_8\Phi_{10}$	1	.	.	1		
1.31^2	$q^{12}\Phi_3^2\Phi_6\Phi_8\Phi_{14}$.	.	.	1	1	.	1	1		
.321	$\frac{1}{2}q^{14}\Phi_2^4\Phi_6\Phi_{10}\Phi_{14}$.	.	.	1	.	.	.	1		
$1.2^2.1$	$\frac{1}{2}q^{14}\Phi_3\Phi_5\Phi_8\Phi_{14}$.	.	1	.	.	1	.	a		
$1^5.1$	$\frac{1}{2}q^{14}\Phi_3\Phi_8\Phi_{10}\Phi_{14}$	1	.	.	.		
1.1^5	$q^{28}\Phi_3\Phi_6$	1	.	.	.		
		$ps \ ^2D_2 \ ps \ ps \ ps \ ps \ ^2D_2 \ ^2D_2 \ ^2D_2 \ ^2D_2 \ .2^2 \ ^2D_2 \ A_3 \ .1^4$									

Theorem 7.5. *Let $(q, 6) = 1$. The decomposition matrices for the unipotent ℓ -blocks of ${}^2E_6(q)$ of positive defect, where $(q^2 + 1)_\ell > 5$, are as given in Tables 26 and 27.*

summands in the principal block of the Harish-Chandra induction from proper Levi subgroups in that series, viz. $\Psi_8, \Psi_9, \Psi_{10}, \Psi_{11}, \Psi_{13}$ and Ψ_{14} . Two further PIMs in that series lie in the block of cyclic defect. This accounts for all non-cuspidal Harish-Chandra series. Hence the four missing columns must correspond to cuspidal Brauer characters.

For the remaining columns we consider the following three GGGRs, whose existence is given by [5, Thm. 6.5(ii)], assuming that p is good:

- the GGGR associated to the family containing ${}^2E_6[1]$ and with projection ${}^2E_6[1] + 2\phi'_{6,6} + \phi_{12,4}$ on this family;
- the GGGR associated to the family $\{\phi'_{2,16}, \phi_{9,10}, \phi''_{1,12}, \phi''_{8,9}\}$ and with projection $\phi''_{1,12} + \phi''_{8,9}$ on this family;
- the GGGR associated to the family $\{\phi''_{2,16}\}$.

From (GGGR) we deduce the unitriangularity of the decomposition matrix. In addition, if we denote by c_1, \dots, c_9 the unknown entries in the 7th column then $c_2 = c_3 = 0$ and $c_1 \leq 2$. Similarly, if d_1, \dots, d_4 denote the entries in the 12th column then $d_1 = 0$ and $d_2 \leq 1$. The last unknown entry (in the 15th column) will be denoted by d_5 .

A Sylow Φ_4 -torus of G has a regular ℓ -character θ whenever $(q^2 + 1)_\ell > 5$. By (Red), the ℓ -reduction of a non-unipotent character induced from θ yields relations on the c_i 's and d_i 's, namely $d_5 \geq 2$, $3 + 2d_2 - 2d_3 + d_4 \geq 0$ and $-4 - 2c_1 - 3c_4 + 3c_5 - c_6 + 2c_7 - 2c_8 + c_9 \geq 0$. To obtain the opposite inequalities we use (DL) successively for the elements $s_1 s_2 s_3 s_4$, $s_1 s_2 s_3 s_1 s_4 s_3$ and $s_1 s_2 s_4 s_3 s_1 s_5 s_4 s_3 s_6 s_5 s_4 s_3$. \square

As in Proposition 3.8, we can use [8, Conj. 1.2] to obtained conjectural upper bounds on the unknown entries in the decomposition matrix which do not depend on q .

Proposition 7.6. *Assume Conjecture 1.2 in [8] holds. Then in the decomposition matrix of the principal Φ_4 -block of ${}^2E_6(q)$, we have $c_1 = 0$, $c_4 \leq 3$, $c_5 \leq 26$, $c_6 \leq 29$, $c_7 \leq 50$, $c_8 \leq 156$ and $d_3 \leq 6$.*

Proof. We consider virtual characters Q_w afforded by the Alvis–Curtis dual of the intersection cohomology of suitably chosen Deligne–Lusztig varieties. In the following table, we give, for each element w we consider, the multiplicity of a PIM Ψ_i in $Q_w[\lambda]$. In order to simplify notation, we denote s_1, \dots, s_6 by $1, \dots, 6$.

w	λ	i	$\langle Q_w[\lambda], \varphi_i \rangle$
1231454236542314356	1	8	$-36c_1$
123423145431	1	11	$3 - c_4$
		12	$23 + c_4 - c_5$
2342314354316543	1	13	$6(29 - c_6)$
		14	$6(50 + c_1 - d_2(23 + c_4 - c_5) - c_7)$
23143154316543	1	15	$2(106 - c_1 + c_7 + (d_2 - d_3)(23 + c_4 - c_5) - c_8)$
546542	-1	15	$5 + d_2 - d_3$

Conjecture 1.2 in [8] predicts that the entries in the last column of the previous table are non-negative. We deduce that $c_1 = 0$, $c_4 \leq 3$, $c_5 \leq c_4 + 23 \leq 26$, $c_6 \leq 29$, $c_7 \leq$

$$50 + c_1 - d_2(23 + c_4 - c_5) \leq 50,$$

$$c_8 \leq 106 - c_1 + c_7 + (d_2 - d_3)(23 + c_4 - c_5) \leq 106 - c_1 + c_7 + d_2(23 + c_4 - c_5) \leq 156$$

and $d_3 \leq d_2 + 5 \leq 6$ since $d_2 \in \{0, 1\}$. \square

7.4. The Φ_4 -blocks in twisted groups. As in the untwisted case we collect some data on the Φ_4 -blocks studied above in Table 28.

TABLE 28. HC-series in Φ_4 -blocks of twisted groups

G	$W_G(b)$	$ \text{IBr}(b) $	ps	2D_2	A_3	$.2^2$	$.1^4$	c
2D_5	$G(4, 1, 2)$	14	5	6	1	1	1	
2D_7			5	6	1	1	1	
2E_6	G_8	16	6	6				4

Note that the decomposition matrices for the blocks in 2D_5 and in 2D_7 of defect Φ_4^2 coincide after permuting rows and columns suitably. Again, it would be interesting to see whether this is caused by a Morita equivalence between these blocks. On the other hand, the multisets of entries of the matrices for twisted groups differ from those for any of the untwisted ones, so if there exists a Morita equivalence between blocks for twisted and untwisted groups, it would have to be with respect to a different choice of basic sets.

8. DECOMPOSITION MATRICES FOR SYMPLECTIC GROUPS AND $F_4(q)$

We now turn to groups with non-simply laced Dynkin diagram, where we start by giving (approximations) to decomposition matrices for the unipotent blocks of small rank symplectic groups $\text{Sp}_{2n}(q)$ for primes $\ell \mid (q^2 + 1)$. Again, it is not known a priori in our present situation that the decomposition matrix has triangular shape. This leads to additional complications.

For completeness and for use in the subsequent proofs, we recall the known Brauer trees for $\text{Sp}_4(q)$ and $\text{Sp}_6(q)$ (see [10]). We also indicate the modular Harish-Chandra series of the PIMs.

Theorem 8.1 (Fong–Srinivasan). *Let $2 \neq \ell \mid (q^2 + 1)$ be a prime. Then the Brauer trees for the unipotent ℓ -blocks of $\text{Sp}_4(q)$ and $\text{Sp}_6(q)$ are as given in Table 29.*

Next, let $G = \text{Sp}_8(q)$.

Theorem 8.2. *Let ℓ be a prime. Assume that q is odd. Then the decomposition matrices for the unipotent ℓ -blocks of $\text{Sp}_8(q)$, $(q^2 + 1)_\ell > 5$, are as given in Tables 30 and 31.*

Proof. The group $G = \text{Sp}_8(q)$ has three unipotent blocks of Φ_4 -defect zero, two blocks of Φ_4 -defect 1 with four characters each, and all other unipotent characters lie in the principal block. The Brauer trees for the blocks of defect 1 are known by [10] and in any case can easily be recovered by (HCi).

The projective modules Ψ_i for $i \in \{1, \dots, 9, 11, 13\}$ are obtained by (HCi). Using (HCr), we can check that there are indecomposable. Finally, (St) yields the last column, leaving

TABLE 29. $\mathrm{Sp}_4(q)$ and $\mathrm{Sp}_6(q)$, $2 \neq \ell | (q^2 + 1)$

$$\begin{array}{llllllll} \mathrm{Sp}_4(q) : & 2. & \text{---} & 1.1 & \text{---} & .1^2 & \text{---} & \bigcirc & \text{---} & C_2 \\ & & ps & & ps & & c & & C_2 & \\ \\ \mathrm{Sp}_6(q) : & 3. & \text{---} & 1.2 & \text{---} & .21 & \text{---} & \bigcirc & \text{---} & C_2 : 1^2 \\ & & & & & & & & & \\ & 21. & \text{---} & 1^2.1 & \text{---} & .1^3 & \text{---} & \bigcirc & \text{---} & C_2 : 2 \\ & & ps & & ps & & .1^2 & & C_2 & \end{array}$$

TABLE 30. $\mathrm{Sp}_8(q)$, $(q^2 + 1)_\ell > 5$

[illegible]

Here, $a \in \{0, 1\}$ and $b \in \{0, 1, 2, 3\}$.

TABLE 31. $\mathrm{Sp}_8(q)$, blocks of defect 1, $2 \neq \ell | (q^2 + 1)$

$$\begin{array}{ccccccc} 3.1 & \text{---} & 2.2 & \text{---} & .2^2 & \text{---} & \bigcirc \text{---} C_2 : .1^2 \\ 2^2. & \text{---} & 1^2.1^2 & \text{---} & 1.1^3 & \text{---} & \bigcirc \text{---} C_2 : 2. \\ & & ps & & ps & & C_2 \end{array}$$

only two (necessarily cuspidal) PIMs to determine. A first approximation of these columns is given by (GGGR) for the families $\{\rho_{1^3,1}, \rho_{1^2,1^2}, \rho_{2,2}, \rho_{C_2,2}\}$ and $\{\rho_{1^4}, \rho_{1,1^3}, \rho_{2,1^2}, \rho_{C_2,1^2}\}$.

We deduce that the relevant submatrix for the last five projectives now has the form

$$\begin{array}{c|ccc}
 C_2 : .2 & 1 & & \\
 1^3.1 & a_1 & 1 & \\
 1^4. & a_2 & 1 & 1 \\
 .21^2 & a_3 & . & a_5 & 1 \\
 .1^4 & a_4 & 1 & a_6 & 1 & 1
 \end{array}$$

In addition, [5, Thm. 6.5(ii)] yields $a_1 \in \{0, 1\}$ and $a_5 = 0$.

As usual, relations on the a_i 's are obtained by looking at suitable Deligne–Lusztig characters: from (Cox) we deduce that $a_1 - a_2 \geq 0$ and $a_1 + a_3 - a_4 + 2 \geq a_6(a_1 - a_2)$. But from (Red) with the induction of an ℓ -character of a Sylow Φ_4 -torus in general position, which exists whenever $(q^2 + 1)_\ell > 5$, we get $a_1 + a_3 - a_4 + 2 \leq a_2 - a_1$. Consequently, $0 \leq a_6(a_1 - a_2) \leq a_2 - a_1 \leq 0$ which forces $a_1 = a_2$ and $a_4 = a_1 + a_3 + 2$. With (DL) applied to the element $w = s_1 s_2 s_3 s_2 s_1 s_2 s_3 s_4$ of the Weyl group we obtain $a_6 \leq 3$.

Finally, we use the GGGR of $\mathrm{Sp}_{10}(q)$ associated to the family $\{\rho_{1^4.1}, \rho_{1^2.1^3}, \rho_{.2^2.1}, \rho_{C_2:.21}\}$ with projection $\rho_{C_2:.21} + \rho_{1^4.1}$ to this family. Cut by the block containing these two characters, the unipotent part of this GGGR is of the form $\rho_{C_2:.21} + \rho_{1^4.1} + \alpha \rho_{.1^5}$. Its Harish-Chandra restriction to G forces $a_3 = 0$. Setting $a = a_1$ and $b = a_6$, we obtain the decomposition matrix as shown in Table 30 \square

Remark 8.3. Kawanaka's conjecture [17, Conj. 2.4.5] would imply $a = 0$ (see also Remark 3.7).

Theorem 8.4. *Let ℓ be a prime. The decomposition matrix for the principal ℓ -block of $F_4(q)$, $(q, 6) = 1$, $(q^2 + 1)_\ell > 5$, is as given in Table 32.*

For the unipotent blocks with cyclic defect, see [18, Lemma 5.4].

Proof. We start from the approximation to the decomposition matrix which was obtained in the thesis of Köhler [18]. The relevant submatrix for the last eight projectives has the following form:

$$\begin{array}{c|cccccc}
 F_4^{II}[1] & 1 & & & & & \\
 F_4^I[1] & . & 1 & & & & \\
 F_4[-I] & . & . & 1 & & & \\
 F_4[I] & . & . & . & 1 & & \\
 \phi_{9,10} & . & . & c_1 & c_1 & 1 & \\
 \phi_{4,13} & a_1 & . & c_2 & c_2 & 1 & 1 \\
 B_2 : .1^2 & . & b_1 & c_3 & c_3 & . & . & 1 \\
 \phi_{1,24} & a_2 & b_2 & c_4 & c_4 & 1 & d & e & 1
 \end{array}$$

First relations come from the ℓ -reduction of non-unipotent characters. For each uniform character $\rho \in \mathcal{E}(G, (s))$ we construct, we give in the table below the type of $C_{\mathbf{G}}(s)$, the Jordan correspondent ρ_s of ρ and the relations that we will use:

$C_{\mathbf{G}}(s)$	ρ_s	relations
$B_2 \cdot (q^2 + 1)$	$\rho_{1^2} + \rho_{B_2}$	$c_2 \geq c_1 - 1$
$(q^2 + 1)^2$	1	$b_2 \geq 2b_1 - 3$
		$c_4 \geq 3c_1 - 2c_2 + 2c_3 - 4$
		$e \geq 2$

TABLE 32. $F_4(q)$, $(q, 6) = 1$, $(q^2 + 1)_\ell > 5$ [illegible]

Here, $a_1 \leq 5$, $a_2 \leq 13 + (5 - a_1)d$, $c_3 \in \{0, 1\}$, $c_4 = c_1 + 2c_3 - 2$ and $d \in \{0, 1, 2\}$.

Now we apply (DL) successively to obtain relations on the unknown entries. Starting with the Deligne–Lusztig character associated with a Coxeter element w we find the non-negative coefficients $c_1 - c_2 - 1$, $3 - 2c_3$ and $2 + 2c_1 - 2c_4 - 3e + 2c_3e$ in R_w of the PIMs corresponding to columns 9, 10, 14 and 16, so that $c_2 = c_1 - 1$ and $c_3 \in \{0, 1\}$. The second relation can then be written $-2(4 - 3c_1 + 2c_2 - 2c_3 + c_4) + (3 - 2c_3)(2 - e) \geq 0$. Since it is a sum of two nonpositive integers, we obtain $4 - 3c_1 + 2c_2 - 2c_3 + c_4 = 0$ and $(3 - 2c_3)(2 - e) = 0$, so that $c_4 = c_1 + 2c_3 - 2$ and $e = 2$. Consequently, the PIMs corresponding to the columns 9, 10, 14 and 16 do not occur in R_w .

We continue with the Deligne–Lusztig character associated with $w' = s_1 s_2 s_3 s_4 s_1 s_2 s_3 s_4$. Using (DL) we find $2b_1 - b_2 - 3 \geq 0$ which forces $b_2 = 2b_1 - 3$ by the previous inequalities. In addition, the PIMs corresponding to the columns 9, 14 and 16 still do not occur in $R_{w'}$.

Finally, with $w'' = s_1 s_2 s_3 s_4 s_1 s_2 s_3 s_4 s_1 s_2 s_3 s_4$ we consider the characters $R_{w''}[\lambda]$ for various eigenvalues λ of F . The multiplicities of the 14th and 16th PIM in these virtual characters yield the relations

$$a_1 \leq 5, \quad d \leq 2, \quad a_2 \leq 13 + (a_1 - 5)d$$

(in particular $a_2 \leq 13$).

Up to finitely many possibilities, the decomposition matrix given above depends only on two unknown parameters, viz. b_1, c_1 . Moreover, these could be bounded above by suitable polynomials in q using GGGRs. As in Proposition 3.8, we can also produce conjectural bounds independent of q .

Proposition 8.5. *Assume Conjecture 1.2 in [8] holds. Then in the decomposition matrix of the principal Φ_4 -block of $F_4(q)$ we have $b_1 \in \{2, 3, 4\}$ and $c_1 \in \{1, 2\}$.*

Proof. We compute the Alvis–Curtis dual of the intersection cohomology of two Deligne–Lusztig varieties, corresponding to the elements $w_1 = s_1 s_2 s_3 s_4$ and $w_2 = s_2 s_4 s_3 s_2 s_1 s_3 s_2 s_3$. Conjecture 1.2 in [8] predicts that the corresponding characters, denoted in [8] by Q_{w_1} and Q_{w_2} , are, up to sign, the unipotent part of projective characters. The same holds for the generalized eigenspaces of F on these characters. The multiplicity of Ψ_{13} in Q_{w_1} is $5 - 2c_1$, which forces $c_1 \leq 2$, and the multiplicity of Ψ_{15} in the 1-eigenspace of F on Q_{w_2} is $4 - b_1$, which forces $b_1 \leq 4$. \square

We collect information on the Harish-Chandra series for the blocks considered in this section in the subsequent Table 33.

TABLE 33. HC-series in Φ_4 -blocks of defect Φ_4^2

G	$W_G(b)$	$ \mathrm{IBr}(b) $	ps	B_2	$.1^2$	C_2	$.1^2$	A_3	c
B_4	$G(4, 1, 2)$	14	6	2	2			1	3
C_4			6			2	2	1	3
F_4	G_8	16	5	1	1	1	1		7

REFERENCES

- [1] K. BREMKE, The decomposition numbers of Hecke algebras of type F_4 with unequal parameters. *Manuscripta Math.* **83** (1994), 331–346.
- [2] M. BROUÉ, Isométries de caractères et équivalences de Morita ou dérivées. *Inst. Hautes Études Sci. Publ. Math.* No. **71** (1990), 45–63.
- [3] M. BROUÉ, G. MALLE, J. MICHEL, Generic blocks of finite reductive groups. *Astérisque* **212** (1993), 7–92.
- [4] P. DELIGNE AND G. LUSZTIG, Representations of reductive groups over finite fields. *Ann. of Math.* **103** (1976), 103–161.
- [5] F. DIGNE, G. LEHRER, J. MICHEL, On character sheaves and characters of reductive groups at unipotent classes. Preprint, arXiv:1307.0698, 2013.
- [6] R. DIPPER, On quotient of Hom-functors and representations of finite general linear groups I. *J. Algebra* **130** (1990), 235–259.
- [7] O. DUDAS, G. MALLE, Decomposition matrices for unitary groups. Submitted, 2013.
- [8] O. DUDAS, G. MALLE, Projective modules in the intersection cohomology of Deligne–Lusztig varieties. *C. R. Acad. Sci. Paris Sér. I Math.* **352** (2014), 467–471.
- [9] O. DUDAS, A note on decomposition numbers for groups of Lie type of small rank. *J. Algebra* **388** (2013), 364–373.
- [10] P. FONG, B. SRINIVASAN, Brauer trees in classical groups. *J. Algebra* **131** (1990), 179–225.
- [11] M. GECK, *Verallgemeinerte Gelfand–Graev Charaktere und Zerlegungszahlen endlicher Gruppen vom Lie-Typ*. Dissertation, RWTH Aachen, 1990.
- [12] M. GECK, G. HISS, Modular representations of finite groups of Lie type in non-defining characteristic. In: *Finite reductive groups: related structures and representations* (ed. M. Cabanes), pp. 195–249, Birkhäuser, Basel, 1997.
- [13] M. GECK, G. HISS, G. MALLE, Cuspidal unipotent Brauer characters. *J. Algebra* **168** (1994), 182–220.

- [14] M. GECK, N. JACON, *Representations of Hecke algebras at roots of unity*. Algebra and Applications, 15. Springer-Verlag, London, 2011.
- [15] M. GECK, J. MÜLLER, James' conjecture for Hecke algebras of exceptional type. I. J. Algebra **321** (2009), 3274–3298.
- [16] M. GECK, G. PFEIFFER, Unipotent characters of the Chevalley groups $D_4(q)$, q odd. Manuscripta Math. **76** (1992), 281–304.
- [17] N. KAWANAKA, Shintani Lifting and Gelfand-Graev Representations. *The Arcata Conference on Representations of Finite Groups* (Arcata, Calif., 1986), 147–163, Proc. Sympos. Pure Math., 47, Part 1. Amer. Math. Soc., Providence, RI, 1987.
- [18] C. KÖHLER, *Unipotente Charaktere und Zerlegungszahlen der endlichen Chevalleygruppen vom Typ F_4* . Dissertation, RWTH Aachen, 2006.
- [19] G. LUSZTIG, *Characters of reductive groups over a finite field*. Annals of Mathematics Studies, 107. Princeton University Press, Princeton, NJ, 1984.
- [20] G. LUSZTIG, A unipotent support for irreducible representations. Adv. Math. **94** (1992), 139–179.
- [21] J. MICHEL, The development version of the CHEVIE package of GAP3. Preprint, arXiv: 1310.7905, 2013.
- [22] H. MIYACHI, Rouquier blocks in Chevalley groups of type E . Adv. Math. **217** (2008), 2841–2871.
- [23] J. TAYLOR, Generalised Gelfand-Graev representations in small characteristics. Preprint, arXiv: 1408.1643, 2014.

UNIVERSITÉ PARIS DIDEROT, UFR DE MATHÉMATIQUES, BÂTIMENT SOPHIE GERMAIN, 5 RUE THOMAS MANN, 75205 PARIS CEDEX 13, FRANCE.

E-mail address: olivier.dudas@imj-prg.fr

FB MATHEMATIK, TU KAISERSLAUTERN, POSTFACH 3049, 67653 KAISERSLAUTERN, GERMANY.

E-mail address: malle@mathematik.uni-kl.de